

North Inlet & Winyah Bay Literature References

Geology & General Ecology

- Dame R, Bushek D, Allen D, Edwards D, Gregory L, Lewitus A, Crawford S, Koepfler E, Corbett C, Kjerfve B, Prins T (2000) The experimental analysis of tidal creeks dominated by oyster reefs: The premanipulation year. *Journal of Shellfish Research* 19:361-369
- Dame R, Childers D, Koepfler E (1992) A Geohydrologic Continuum Theory for the Spatial and Temporal Evolution of Marsh-Estuarine Ecosystems. *Netherlands Journal of Sea Research* 30:63-72
- Dame R, Chrzanowski T, Bildstein K, Kjerfve B, McKellar H, Nelson D, Spurrier J, Stancyk S, Stevenson H, Vernberg J, Zingmark R (1986) The Outwelling Hypothesis and North Inlet, South-Carolina. *Marine Ecology-Progress Series* 33:217-229
- Dame RF, Gardner LR (1993) Nutrient Processing and the Development of Tidal Creek Ecosystems. *Marine Chemistry* 43:175-183
- Dame RF, Spurrier JD, Williams TM, Kjerfve B, Zingmark RG, Wolaver TG, Chrzanowski TH, McKellar HN, Vernberg FJ (1991) Annual Material Processing by a Salt-Marsh Estuarine-Basin in South-Carolina, USA. *Marine Ecology-Progress Series* 72:153-166
- Gardner LR, Bohn M (1980) Geomorphic and Hydraulic Evolution of Tidal Creeks on a Subsiding Beach Ridge Plain, North-Inlet, Sc. *Marine Geology* 34:M91-M97
- Gardner LR, Michener WK, Williams TM, Blood ER, Kjerfve B, Smock LA, Lipscomb DJ, Gresham C (1992) Disturbance Effects of Hurricane Hugo on a Pristine Coastal Landscape - North Inlet, South-Carolina, USA. *Netherlands Journal of Sea Research* 30:249-263
- Gardner LR (2005) Role of geomorphic and hydraulic parameters in governing pore water seepage from salt marsh sediments. *Water Resources Research* 41
- Novakowski KI, Torres R, Gardner LR, Voulgaris G (2004) Geomorphic analysis of tidal creek networks. *Water Resources Research* 40

Hydrology

- Carter ES, White SM, Wilson AM (2008) Variation in groundwater salinity in a tidal salt marsh basin, North Inlet Estuary, South Carolina. *Estuarine Coastal and Shelf Science* 76:543-552
- Gardner LR, Reeves HW (2002) Seasonal patterns in the soil water balance of a *Spartina* marsh site at North Inlet, South Carolina, USA. *Wetlands* 22:467-477
- Gardner LR, Wilson AM (2006) Comparison of four numerical models for simulating seepage from salt marsh sediments. *Estuarine Coastal and Shelf Science* 69:427-437
- Hutchinson SE, Sklar FH (1993) Lunar Periods as Grouping Variables for Temporally Fixed Sampling Regimes in a Tidally Dominated Estuary. *Estuaries* 16:789-798
- Kjerfve B, Miranda LB, Wolanski E (1991) Modeling Water Circulation in an Estuary and Intertidal Salt-Marsh System. *Netherlands Journal of Sea Research* 28:141-147
- Miller JL, Gardner LR (1981) Sheet Flow in a Salt-Marsh Basin, North Inlet, South-Carolina. *Estuaries* 4:234-237
- Morris JT (1995) The Mass-Balance of Salt and Water in Intertidal Sediments - Results from North-Inlet, South-Carolina. *Estuaries* 18:556-567
- Traynum S, Styles R (2008) Exchange flow between two estuaries connected by a shallow tidal channel. *Journal of Coastal Research* 24:1260-1268
- Wilson AM, Gardner LR (2006) Tidally driven groundwater flow and solute exchange in a marsh: Numerical simulations. *Water Resources Research* 42

Uncles RJ, Kjerfve B (1986) Transverse Structure of Residual Flow in North Inlet, South-Carolina. *Estuaries* 9:39-42

Sediment dynamics

Gardner LR, Kjerfve B (2006) Tidal fluxes of nutrients and suspended sediments at the North Inlet - Winyah Bay National Estuarine Research Reserve. *Estuarine Coastal and Shelf Science* 70:682-692

Gardner LR, Thombs L, Edwards D, Nelson D (1989) Time-Series Analyses of Suspended Sediment Concentrations at North Inlet, South-Carolina. *Estuaries* 12:211-221

Hutchinson SE, Sklar FH, Roberts C (1995) Short-Term Sediment Dynamics in a Southeastern USA *Spartina* Marsh. *Journal of Coastal Research* 11:370-380

Murphy S, Voulgaris G (2006) Identifying the role of tides, rainfall and seasonality in marsh sedimentation using long-term suspended sediment concentration data. *Marine Geology* 227:31-50

Mwamba MJ, Torres R (2002) Rainfall effects on marsh sediment redistribution, North Inlet, South Carolina, USA. *Marine Geology* 189:267-287

Torres R, Mwamba MJ, Goni MA (2003) Properties of intertidal marsh sediment mobilized by rainfall. *Limnology and Oceanography* 48:1245-1253

Traynum S, Styles R (2007) Flow, stress and sediment resuspension in a shallow tidal channel. *Estuaries and Coasts* 30:94-101

Voulgaris G, Meyers ST (2004) Net effect of rainfall activity on salt-marsh sediment distribution. *Marine Geology* 207:115-129

Wargo CA, Styles R (2007) Along channel flow and sediment dynamics at North Inlet, South Carolina. *Estuarine Coastal and Shelf Science* 71:669-682

Wolaver TG, Dame RF, Spurrier JD, Miller AB (1988) Bly Creek Ecosystem Study - Inorganic Sediment Transport within an Euhaline Salt-Marsh Basin, North Inlet, South-Carolina. *Journal of Coastal Research* 4:607-615

Carbon, Nutrient & Water Quality

Apple JK, Smith EM, Boyd TJ (2008) Temperature, salinity, nutrients and the covariation of bacterial production and chlorophyll a in estuarine ecosystems. *Journal of Coastal Research* 55: 59-75

Buzzelli C, Akman O, Buck T, Koepfler E, Morris J, Lewitus A (2004) Relationships among water-quality parameters from the North Inlet-Winyah Bay National Estuarine Research Reserve, South Carolina. *Journal of Coastal Research*:59-74

Cai WJ, Wang YC, Krest J, Moore WS (2003) The geochemistry of dissolved inorganic carbon in a surficial groundwater aquifer in North Inlet, South Carolina, and the carbon fluxes to the coastal ocean. *Geochimica Et Cosmochimica Acta* 67:631-639

Childers DL, McKellar HN, Dame RF, Sklar FH, Blood ER (1993) A Dynamic Nutrient Budget of Subsystem Interactions in a Salt-Marsh Estuary. *Estuarine Coastal and Shelf Science* 36:105-131

Chrzanowski TH, Stevenson LH, Spurrier JD (1982) Transport of Particulate Organic-Carbon through the North Inlet Ecosystem. *Marine Ecology-Progress Series* 7:231-245

Chrzanowski TH, Stevenson LH, Spurrier JD (1983) Transport of Dissolved Organic-Carbon through a Major Creek of the North Inlet Ecosystem. *Marine Ecology-Progress Series* 13:167-174

- Dame RF, Stilwell D (1984) Environmental-Factors Influencing Macrodetritus Flux in North Inlet Estuary. *Estuarine Coastal and Shelf Science* 18:721-726
- Dame RF, Wolaver TG, Williams TM, Spurrier JD, Miller AB (1990) The Bly Creek Ecosystem Study - Phosphorus Transport within a Euhaline Salt-Marsh Basin, North Inlet, South-Carolina. *Netherlands Journal of Sea Research* 27:73-80
- D'Andrea AF, Aller RC, Lopez GR (2002) Organic matter flux and reactivity on a South Carolina sandflat: The impacts of porewater advection and macrobiological structures. *Limnology and Oceanography* 47:1056-1070
- Fulton MH, Scott GI, Fortner A, Bidleman TF, Ngabe B (1993) The Effects of Urbanization on Small High Salinity Estuaries of the Southeastern United-States. *Archives of Environmental Contamination and Toxicology* 25:476-484
- Gardner LR, Kjerfve B (2006) Tidal fluxes of nutrients and suspended sediments at the North Inlet - Winyah Bay National Estuarine Research Reserve. *Estuarine Coastal and Shelf Science* 70:682-692
- Gardner LR, Kjerfve B, Petrecca DM (2006) Tidal fluxes of dissolved oxygen at the North Inlet-Winyah Bay National Estuarine Research Reserve. *Estuarine Coastal and Shelf Science* 67:450-460
- Gardner LR, Wolaver TG, Mitchell M (1988) Spatial Variations in the Sulfur Chemistry of Salt-Marsh Sediments at North-Inlet, South-Carolina. *Journal of Marine Research* 46:815-836
- Goni MA, Gardner LR (2003) Seasonal dynamics in dissolved organic carbon concentrations in a coastal water-table aquifer at the forest-marsh interface. *Aquatic Geochemistry* 9:209-232
- Goni MA, Thomas KA (2000) Sources and transformations of organic matter in surface soils and sediments from a tidal estuary (north inlet, South Carolina, USA). *Estuaries* 23:548-564
- Goni MA, Teixeira MJ, Perkey DW (2003) Sources and distribution of organic matter in a river-dominated estuary (Winyah Bay, SC, USA). *Estuarine Coastal and Shelf Science* 57:1023-1048
- Guentzel JL, Tsukamoto Y (2001) Processes influencing mercury speciation and bioconcentration in the North Inlet-Winyah Bay Estuary, South Carolina, USA. *Marine Pollution Bulletin* 42:615-619
- Haertel-Borer SS, Allen DM, Dame RF (2004) Fishes and shrimps are significant sources of dissolved inorganic nutrients in intertidal salt marsh creeks. *Journal of Experimental Marine Biology and Ecology* 311:79-99
- Huang XQ, Morris JT (2005) Distribution of phosphatase activity in marsh sediments along an estuarine salinity gradient. *Marine Ecology-Progress Series* 292:75-83
- Krest JM, Moore WS, Gardner LR, Morris JT (2000) Marsh nutrient export supplied by groundwater discharge: Evidence from radium measurements. *Global Biogeochemical Cycles* 14:167-176
- Kucklick JR, Bidleman TF (1994) Organic Contaminants in Winyah Bay, South-Carolina .1. Pesticides and Polycyclic Aromatic-Hydrocarbons in Subsurface and Microlayer Waters. *Marine Environmental Research* 37:63-78
- Nelson KA, Scott GI, Rust PF (2005) A multivariable approach for evaluating major impacts on water quality in Murrells and North Inlets, south Carolina. *Journal of Shellfish Research* 24:1241-1251
- Paludan C, Morris JT (1999) Distribution and speciation of phosphorus along a salinity gradient in intertidal marsh sediments. *Biogeochemistry* 45:197-221

- White DL, Porter DE, Lewitus AJ (2004) Spatial and temporal analyses of water quality and phytoplankton biomass in an urbanized versus a relatively pristine salt marsh estuary. *Journal of Experimental Marine Biology and Ecology* 298:255-273
- Williams TM, Wolaver TG, Dame RF, Spurrier JD (1992) The Bly Creek Ecosystem Study - Organic-Carbon Transport within a Euhaline Salt-Marsh Basin, North Inlet, South-Carolina. *Journal of Experimental Marine Biology and Ecology* 163:125-139
- Wolaver TG, Gardner LR (1983) Salt-Marsh Sediment Interstitial Iron and Sulfate Concentrations Along a Height Gradient from Tidal Creeks to the High Marsh in North Inlet, South-Carolina. *Estuaries* 6:296-297
- Wolaver TG, Hutchinson S, Marozas M (1986) Dissolved and Particulate Organic-Carbon in the North Inlet Estuary, South-Carolina - What Controls Their Concentrations. *Estuaries* 9:31-38
- Wolaver TG, Johnson W, Marozas M (1984) Nitrogen and Phosphorus Concentrations within North Inlet, South-Carolina - Speculation as to Sources and Sinks. *Estuarine Coastal and Shelf Science* 19:243-255
- Wolaver TG, Whiting GJ, Dame RF, Williams TM, Spurrier JD (1988) Bly Creek Ecosystem Study - Nitrogen Exchange within a Euhaline Salt-Marsh Basin of North Inlet, South-Carolina. *Marine Ecology-Progress Series* 49:107-116

Primary producers

- Dame RF, Kenny PD (1986) Variability of *Spartina-Alterniflora* Primary Production in the Euhaline North Inlet Estuary. *Marine Ecology-Progress Series* 32:71-80
- Flavier AB, Zingmark RG (1993) Macroalgal Recruitment in a High Marsh Creek of North Inlet Estuary, South-Carolina. *Journal of Phycology* 29:2-8
- Jackson G, Zingmark R, Lewitus AJ, Tymowski RG, Stuckey J (2006) Spatial and temporal dynamics of epiphytic microalgae on the cordgrass *Spartina alterniflora* in North Inlet Estuary, South Carolina. *Estuaries and Coasts* 29:1212-1221
- Lewitus AJ, Kawaguchi T, DiTullio GR, Keese JDM (2004) Iron limitation of phytoplankton in an urbanized vs. forested southeastern US salt marsh estuary. *Journal of Experimental Marine Biology and Ecology* 298:233-254
- Lewitus AJ, Koepfler ET, Morris JT (1998) Seasonal variation in the regulation of phytoplankton by nitrogen and grazing in a salt-marsh estuary. *Limnology and Oceanography* 43:636-646
- Lewitus AJ, White DL, Tymowski RG, Geesey ME, Hymel SN, Noble PA (2005) Adapting the CHEMTAX method for assessing phytoplankton taxonomic composition in southeastern US estuaries. *Estuaries* 28:160-172
- Morris JT (2006) Competition among marsh macrophytes by means of geomorphological displacement in the intertidal zone. *Estuarine Coastal and Shelf Science* 69:395-402
- Morris JT, Porter D, Neet M, Noble PA, Schmidt L, Lapine LA, Jensen JR (2005) Integrating LIDAR elevation data, multi-spectral imagery and neural network modelling for marsh characterization. *International Journal of Remote Sensing* 26:5221-5234
- Noble PA, Tymowski RG, Fletcher M, Morris JT, Lewitus AJ (2003) Contrasting patterns of phytoplankton community pigment composition in two salt marsh estuaries in southeastern United States. *Applied and Environmental Microbiology* 69:4129-4143
- Pinckney J, Zingmark RG (1991) Effects of Tidal Stage and Sun Angles on Intertidal Benthic Microalgal Productivity. *Marine Ecology-Progress Series* 76:81-89

- Pinckney J, Zingmark RG (1993) Biomass and Production of Benthic Microalgal Communities in Estuarine Habitats. *Estuaries* 16:887-897
- Pinckney J, Zingmark RG (1993) Photophysiological Responses of Intertidal Benthic Microalgal Communities to in-Situ Light Environments - Methodological Considerations. *Limnology and Oceanography* 38:1373-1383
- Pinckney JL, Zingmark RG (1993) Modeling the Annual Production of Intertidal Benthic Microalgae in Estuarine Ecosystems. *Journal of Phycology* 29:396-407
- Wetz MS, Hayes KC, Lewitus AJ, Wolny JL, White DL (2006) Variability in phytoplankton pigment biomass and taxonomic composition over tidal cycles in a salt marsh estuary. *Marine Ecology-Progress Series* 320:109-120
- White DL, Porter DE, Lewitus AJ (2004) Spatial and temporal analyses of water quality and phytoplankton biomass in an urbanized versus a relatively pristine salt marsh estuary. *Journal of Experimental Marine Biology and Ecology* 298:255-273

Microbes

- Apple JK, Smith EM, Boyd TJ (2008) Temperature, salinity, nutrients and the covariation of bacterial production and chlorophyll a in estuarine ecosystems. *Journal of Coastal Research* 55: 59-75
- DeLorenzo ME, Lewitus AJ, Scott GI, Ross PE (2001) Use of metabolic inhibitors to characterize ecological interactions in an estuarine microbial food web. *Microbial Ecology* 42:317-327
- Bagwell CE, Piceno YM, Ashburne-Lucas A, Lovell CR (1998) Physiological diversity of the rhizosphere diazotroph assemblages of selected salt marsh grasses. *Applied and Environmental Microbiology* 64:4276-4282
- Johnson W, Lewitus AJ, Fletcher M (2006) Linking bacterioplankton community structures to environmental state variables and phytoplankton assemblages in two South Carolina salt marsh estuaries. *Aquatic Microbial Ecology* 45:129-145
- Lovell CR, Friez MJ, Longshore JW, Bagwell CE (2001) Recovery and phylogenetic analysis of nifH sequences from diazotrophic bacteria associated with dead aboveground biomass of *Spartina alterniflora*. *Applied and Environmental Microbiology* 67:5308-5314
- Lovell CR, Piceno YM, Quattro JM, Bagwell CE (2000) Molecular analysis of diazotroph diversity in the rhizosphere of the smooth cordgrass, *Spartina alterniflora*. *Applied and Environmental Microbiology* 66:3814-3822
- Noble PA, Almeida JS, Lovell CR (2000) Application of neural computing methods for interpreting phospholipid fatty acid profiles of natural microbial communities. *Applied and Environmental Microbiology* 66:694-699
- Wetz MS, Lewitus AJ, Koepfler ET, Hayes KC (2002) Impact of the Eastern oyster *Crassostrea virginica* on microbial community structure in a salt marsh estuary. *Aquatic Microbial Ecology* 28:87-97
- Chrzanowski TH, Stevenson LH, Spurrier JD (1982) Transport of Microbial Biomass through the North Inlet Ecosystem. *Microbial Ecology* 8:139-156

Benthic fauna

- Brodie RJ, Behum ME, Monroe E, Glenn N, Staton JL (2005) Recruitment to adult habitats following marine planktonic development in the fiddler crabs, *Uca pugnator*, *U. pugnax*, and *U. minax*. *Marine Biology* 147:105-111

- Dame RF, Spurrier JD, Zingmark RG (1992) In situ Metabolism of an Oyster Reef. *Journal of Experimental Marine Biology and Ecology* 164:147-159
- Feller RJ, Stancyk SE, Coull BC, Edwards DG (1992) Recruitment of Polychaetes and Bivalves - Long-Term Assessment of Predictability in a Soft-Bottom Habitat. *Marine Ecology-Progress Series* 87:227-238
- Godley J, Brodie R (2007) Effect of summer storms on early life stages of *Uca minax*, *U. pugnax* and *U. pugilator* in North Inlet Estuary, South Carolina, USA. *Marine Ecology-Progress Series* 342:197-204
- Hartman MJ, Stancyk SE (2000) Distribution of an invasive anomuran decapod, *Petrolisthes armatus*, in the north inlet-Winyah Bay National Estuarine Research Reserve on the South Carolina coast. *American Zoologist* 40:1047-1047
- Hartman MJ, Stancyk SE, Lohrer AM (2002) Native predators and the success of a newcomer *Petrolisthes armatus*, the green porcelain crab to an oyster reef ecosystem in North Inlet, SC. *Integrative and Comparative Biology* 42:1241-1241
- Michener WK, Kenny PD (1991) Spatial and Temporal Patterns of *Crassostrea-Virginica* (Gmelin) Recruitment - Relationship to Scale and Substratum. *Journal of Experimental Marine Biology and Ecology* 154:97-121
- Service SK, Feller RJ (1992) Long-Term Trends of Subtidal Macrobenthos in North Inlet, South-Carolina. *Hydrobiologia* 231:13-40
- Wetz MS, Lewitus AJ, Koepfler ET, Hayes KC (2002) Impact of the Eastern oyster *Crassostrea virginica* on microbial community structure in a salt marsh estuary. *Aquatic Microbial Ecology* 28:87-97

Zooplankton

- Allen DM, Ogburn-Matthews V, Buck T, Smith EM (2008) Mesozooplankton response to climate change and variability in a southeastern US Estuary. *Journal of Coastal Research* 55: 95-110
- Allen DM, Edwards D, Feller RJ, Hutchinson SE, OgburnMatthews V (1997) Detection and analysis of unusual events in long-term zooplankton and nekton data sets from North Inlet Estuary, South Carolina, USA. *Oceanologica Acta* 20:165-175
- Barans CA, Stender BW, Holliday DV, Greenlaw CF (1997) Variation in the vertical distribution of zooplankton and fine particles in an estuarine inlet of South Carolina. *Estuaries* 20:467-482
- Houser DS, Allen DM (1996) Zooplankton dynamics in an intertidal salt-marsh basin. *Estuaries* 19:659-673
- Johnson, W.S. and D.M. Allen. 2005. *Zooplankton of the Atlantic and Gulf coasts: a guide to their identification and ecology*. Johns Hopkins University Press. Baltimore. 379 pp. ISBN 0-8018-8168-4.
- Marshallon D, Pinckney JL (2007) Respiration rates of dominant hydromedusae in the North Inlet tidal estuary during winter and summer. *Journal of Plankton Research* 29:1031-1040
- Marshallon D, Pinckney JL (2008) Grazing and assimilation rate estimates of hydromedusae from a temperate tidal creek system. *Hydrobiologia* 606:203-211
- Morris JT, Coull BC (1992) Population-Dynamics, Numerical Production, and Potential Predation Impact on a Meiobenthic Copepod. *Canadian Journal of Fisheries and Aquatic Sciences* 49:609-616

Stancyk SE, Christy JH (1981) Movement of Larvae from North Inlet, Sc, with Special Reference to Crab Zoeae. *Estuaries* 4:240-240

Nekton

Abel DC, Young RF, Garwood JA (2007) Survey of the shark fauna in two South Carolina estuaries and the impact of salinity structure. *American Fisheries Society Symposium*:109-124

Allen DM, Haertel-Borer SS, Milan BJ, Bushek D, Dames RF (2007) Geomorphological determinants of nekton use of intertidal salt marsh creeks. *Marine Ecology-Progress Series* 329:57-71

Allen DM, Johnson WS, Ogburnmatthews V (1995) Trophic Relationships and Seasonal Utilization of Salt-Marsh Creeks by Zooplanktivorous Fishes. *Environmental Biology of Fishes* 42:37-50

Allen, D.M., S.K. Service, and M.V. Ogburn. 1992. Factors influencing the collection efficiency of estuarine fishes. *Trans. Am. Fish. Soc.* 121:234-244

Allen, D.M. and D.L. Barker. 1990. Interannual variability in larval fish recruitment to estuarine epibenthic habitats. *Mar. Ecol. Prog. Ser.* 63:113-125.

Beseres JJ, Feller RJ (2007) Changes in the spatial distribution of subtidal macrobenthos due to predation by white shrimp (*Litopenaeus setiferus*). *Estuaries and Coasts* 30:591-600

Beseres JJ, Feller RJ (2007) Importance of predation by white shrimp *Litopenaeus setiferus* on estuarine subtidal macrobenthos. *Journal of Experimental Marine Biology and Ecology* 344:193-205

Bretsch K, Allen DM (2006) Tidal migrations of nekton in salt marsh intertidal creeks. *Estuaries and Coasts* 29:474-486

Haertel-Borer SS, Allen DM, Dame RF (2004) Fishes and shrimps are significant sources of dissolved inorganic nutrients in intertidal salt marsh creeks. *Journal of Experimental Marine Biology and Ecology* 311:79-99

Lehnert RL, Allen DM (2002) Nekton use of subtidal oyster shell habitat in a southeastern US estuary. *Estuaries* 25:1015-1024

Ogburn-Matthews, V. and D.M. Allen. 1993. Interactions among some dominant estuarine nekton species. *Estuaries* 16:840-850.

Potthoff MT, Allen DM (2003) Site fidelity, home range, and tidal migrations of juvenile pinfish, *Lagodon rhomboides*, in salt marsh creeks. *Environmental Biology of Fishes* 67:231-240

Reichert MJM (2003) Diet, consumption, and growth of juvenile fringed flounder (*Etropus crossotus*); a test of the 'maximum growth/optimum food hypothesis' in a subtropical nursery area. *Journal of Sea Research* 50:97-116

Vernberg FJ, Piyatiratitivorakul S (1998) Effects of salinity and temperature on the bioenergetics of adult stages of the grass shrimp (*Palaemonetes pugio* Holthuis) from the north inlet estuary, South Carolina. *Estuaries* 21:176-193

Young RF, Phillips HD (2002) Primary production required to support bottlenose dolphins in a salt marsh estuarine creek system. *Marine Mammal Science* 18:358-373