








































Target Species

COMMON PHYTOPLANKTON KEY

OTHER COMMON PLANKTON (non-phyto)

<i>Alexandrium</i> spp. AL  25-46 µm	<i>Gymnodinium</i> spp. GY  24-50 µm	<i>Gonyaulax spinifera</i> GS  25-50 µm	<i>Protoperdinium</i> spp. PT  50-95 µm	<i>Scropsiella</i> spp. SC  20-37 µm	<i>Coccinodiscus</i> spp. CO  40-500 µm	<i>Odontella</i> spp. OD  45-70 µm	<i>Larval Clam</i> LC  Generally Large
<i>Dinophysis norvegica</i> DN  48-80 µm	<i>Dinophysis acuminata</i> DA  40 - 50 µm	<i>Dinophysis tripos</i> DT  40 - 120 µm	<i>Asterionellopsis</i> spp. AS  30-150 µm	<i>Chaetoceros</i> spp. CH  10 - 53 µm	<i>Chaetoceros socialis</i> CS  3-15 µm	<i>Biddulphia</i> spp. BD  60 - 160 µm	<i>Rotifer</i> spp. RO  Generally Large
<i>Prorocentrum lima</i> PL  31-47 µm	<i>Prorocentrum micans</i> PM  35-70 µm	<i>Ceratium fusus</i> CF  200-540 µm	<i>Ceratium lineatum</i> CL  100-130 µm	<i>Ceratium longipes</i> CP  150-250 µm	<i>Dictyocha</i> spp. DO  10-45 µm	<i>Fragilaria</i> spp. FR  10 - 70 µm	<i>Pollen Grain</i> PG  Generally Large
<i>Pseudonitzschia</i> PS  64-117 µm	<i>Thalassionema</i> spp. TA  16 - 90 µm	<i>Thalassiosira</i> spp. TL  12-39 µm	<i>Nitzschia</i> spp. NZ  60 - 125 µm	<i>Skeletonema</i> spp. SK  2-21 µm	<i>Ditylum</i> spp. DM  80 - 130 µm	<i>Leptocylindrus</i> spp. LP  30 - 75 µm	<i>Crab Zoa</i> CZ  Generally Large
<i>Species Name</i> CODE (Guide to using key) illustration of organism Size Range (in µm)	<i>Rhizosolenia</i> spp. RH  25-57 µm	<i>Gyrosigma</i> spp. GY  110 - 175 µm	<i>Navicula</i> spp. NV  32-49 µm	<i>Melosira</i> spp. ML  10-50 µm	<i>Guinardia</i> spp. GN  60 - 160 µm	<i>Eucampia</i> spp. EU  10-33 µm	<i>Tintinnid</i> spp. TN  Generally Large

REFERENCES USED

Cupp, E.E. 1943. Marine Plankton Diatoms of the West Coast of North America, University of California Berkeley
 McConnaughey, B.H. 1970. Introduction to Marine Biology. C.V. Mosby St. Louis
 Smith, D.L., 1997. A Guide to Marine and Coastal Plankton and Marine Invertebrate Larvae; Kendall/Hunt Dubuque.
 Tomas, C.R. 1997. Identifying Marine Phytoplankton. Academic Press/Harcourt Brace San Diego.
 van den Hoek, C., Mann D.G. & Jahns, H.M. 1993. Algae; an introduction to phycology. Cambridge University Press.
<http://www.marbot.gu.s/SSS/SSSHome.htm>

